arXiv:0710.2340v3 [math.AG] 25 Mar 2008

On ground fields of arithmetic hyperbolic reflection groups. III

Viacheslav V. Nikulin*

Abstract

This paper continues [18] (arXiv.org:math.AG/0609256), [19] (arXiv:0708.3991) and [20](arXiv:0710.0162)

Using authors's methods of 1980, 1981, some explicit finite sets of number fields containing all ground fields of arithmetic hyperbolic reflection groups in dimension at least 3 are defined, and explicit bounds of their degrees (over $\mathbb{Q}$) are obtained.

Thus, now, explicit bound of degree of ground fields of arithmetic hyperbolic reflection groups is known in all dimensions. Thus, now, we can, in principle, obtain effective finite classification of arithmetic hyperbolic reflection groups in all dimensions together.

To 85th Birthday of Igor Rostislavich Shafarevich

1 Introduction

This paper continues [18], [19] and [20]. See introductions of these papers about history, definitions and results concerning the subject.

In [19] and [20] some explicit finite sets of totally real algebraic number fields containing all ground fields of arithmetic hyperbolic reflection groups in dimensions $n \geq 4$ were defined, and good explicit bounds of degrees (over $\mathbb{Q}$) of their fields were obtained. In particular, an explicit bound (≤ 56) for $n \geq 6$ and (≤ 138) for $n \geq 4$ of degree of the ground field of any arithmetic hyperbolic reflection group in dimension $n \geq 4$ was obtained.

Here we continue this study for dimensions $n = 3$. Using similar methods, we define some explicit finite sets of totally real algebraic number fields containing all ground fields of arithmetic hyperbolic reflection groups in dimensions $n \geq 3$. Moreover, an explicit bound (≤ 909) of degrees of fields from these sets are obtained. Here we also use result by Long, Maclachlan and Reid from [13].

---

*This paper was written with the financial support of EPSRC, United Kingdom (grant no. EP/D061997/1)

Thus, degree of the ground field of any arithmetic hyperbolic reflection group of dimension $n \geq 3$ is bounded by 909. Remark that finiteness of the number of maximal arithmetic hyperbolic reflection groups of dimension 3 was obtained by Agol [1], it follows a theoretical existence of some bound of the degree of ground fields of arithmetic hyperbolic reflection groups in dimension 3. Difference of our result here is that we give an explicit bound for the degree which is important for finite effective classification. It also gives another proof of finiteness in dimension 3. In fact, using our methods, we show that finiteness in dimension 3 follows from finiteness in dimension 2.

It is also very important that all these fields are attached to fundamental chambers of arithmetic hyperbolic reflection groups, and they can be further geometrically investigated and restricted.

It was also shown in [19] (using results of [13] and [4], [24]) that degree of the ground field of any arithmetic hyperbolic reflection group of dimension $n = 2$ is bounded by 44.

Thus, now an explicit bound of degree of ground fields of arithmetic hyperbolic reflection groups is known in all dimensions. By [14] and [15] and [27], [28], then there exists an effective finite classification of maximal arithmetic hyperbolic reflection groups in all dimensions together. More generally, there exists an effective finite classification of similarity classes of reflective hyperbolic lattices S. Their full reflection groups $W(S)$ contain all maximal arithmetic hyperbolic reflection groups as subgroups of finite index.

Since this paper is a direct continuation of [19] and [20], we use notations, definitions and results of these papers.

2 Ground fields of arithmetic hyperbolic reflection groups in dimensions $n \geq 3$

Since this paper is a direct continuation of [19] and [20], we use notations, definitions and results of this papers.

In [19, Secs 3 and 4], explicit finite sets $\mathcal{F}L^4$, $\mathcal{F}T$, $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 5$, and $\mathcal{F}\Gamma_{2,4}(14)$ of totally real algebraic number fields were defined. The set $\mathcal{F}L^4$ consists of all ground fields of arithmetic Lannér diagrams with ≥ 4 vertices and consists of three fields of degree ≤ 2. The set $\mathcal{F}T$ consists of all ground fields of arithmetic triangles (plane) and has 13 fields of degree ≤ 5 (it includes $\mathcal{F}L^4$). The set $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 5$, consists of all ground fields of V-arithmetic finite edge polyhedra of minimality 14 with connected Gram graph having 4 vertices. They are determined by 5 types of graphs $\Gamma_i^{(4)}(14)$, $i = 1, 2, 3, 4, 5$. The degrees of fields from these sets are bounded by 22, 39, 53, 56, 54 respectively. The set $\mathcal{F}\Gamma_{2,4}(14)$ consists of all ground fields of arithmetic quadrangles (plane) of minimality 14. Their degrees are bounded by 11.

The following result was obtained in [19, Theorem 4.5] using methods of [14] and [15] and results by Borel [4] and Takeuchi [24].

Theorem 2.1. *([19]) In dimensions $n \geq 6$, the ground field of any arithmetic hyperbolic reflection group belongs to one of finite sets of fields $\mathcal{F}L^4$, $\mathcal{F}T$, $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 5$, and $\mathcal{F}\Gamma_{2,4}(14)$. In particular, its degree is bounded by* 56.

In [20, Secs 2 and 3], further explicit finite sets of fields $\mathcal{F}\Gamma_1^{(6)}(14)$, $\mathcal{F}\Gamma_2^{(6)}(14)$, $\mathcal{F}\Gamma_3^{(6)}(14)$, $\mathcal{F}\Gamma_1^{(7)}(14)$, $\mathcal{F}\Gamma_2^{(7)}(14)$, $\mathcal{F}\Gamma_{2,5}(14)$ were defined. The sets $\mathcal{F}\Gamma_1^{(6)}(14)$, $\mathcal{F}\Gamma_2^{(6)}(14)$, $\mathcal{F}\Gamma_3^{(6)}(14)$, $\mathcal{F}\Gamma_1^{(7)}(14)$, $\mathcal{F}\Gamma_2^{(7)}(14)$ are defined by some V-arithmetic pentagon graphs of minimality 14. They are related to some fundamental pentagons on hyperbolic plane. The degree of fields from $\mathcal{F}\Gamma_1^{(6)}(14)$ is bounded by 56, from $\mathcal{F}\Gamma_2^{(6)}(14)$ by 75, from $\mathcal{F}\Gamma_3^{(6)}(14)$ by 138, from $\mathcal{F}\Gamma_1^{(7)}(14)$ by 38, from $\mathcal{F}\Gamma_2^{(7)}(14)$ by 138. The set $\mathcal{F}\Gamma_{2,5}(14)$ consists of all ground fields of arithmetic pentagons (plane) of minimality 14. Their degrees are bounded by 12.

Using similar, but much more complicated considerations, we proved in [20]

Theorem 2.2. *In dimensions $n \geq 4$, the ground field of any arithmetic hyperbolic reflection group belongs to one of finite sets of fields $\mathcal{F}L^4$, $\mathcal{F}T$, $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 5$, $\mathcal{F}\Gamma_{2,4}(14)$ and $\mathcal{F}\Gamma_1^{(6)}(14)$, $\mathcal{F}\Gamma_2^{(6)}(14)$, $\mathcal{F}\Gamma_3^{(6)}(14)$, $\mathcal{F}\Gamma_1^{(7)}(14)$, $\mathcal{F}\Gamma_2^{(7)}(14)$, $\mathcal{F}\Gamma_{2,5}(14)$.*

In particular, its degree is bounded by 138.

Applying the same methods, here we want to extend this result to $n \geq 3$, also considering $n = 3$.

First, we introduce some other explicit finite sets of fields. All of them are related to fundamental polygons on hyperbolic plane.

We consider arithmetic reflection groups on hyperbolic plane with fundamental polygons $\mathcal{M}_2$ of minimality 14. It means that $\delta_1 \cdot \delta_2 < 14$ for any $\delta_1, \delta_2 \in P(\mathcal{M}_2)$. The corresponding polygons $\mathcal{M}_2$ are also called arithmetic polygons of the minimality 14.

Definition 2.3. *We denote by $\Gamma_2(14)$ the set of Gram graphs $\Gamma(P(\mathcal{M}_2))$ of all arithmetic polygons $\mathcal{M}_2$ of minimality* 14. *The set $\mathcal{F}\Gamma_2(14)$ consists of all their ground fields.*

It follows from results of Long, Maclachlan and Reid [13], Borel [4] and Takeuchi [24] (see [19, Sec. 4.5]) that the degree of ground fields of arithmetic hyperbolic reflection groups of dimension two is bounded by 44. Thus, the degree of fields from $\mathcal{F}\Gamma_2(14)$ is also bounded by 44.

Let us consider V-arithmetic 3-dimensional chambers which are defined by the Gram graphs $\Gamma_6^{(4)}$ with 4 vertices $\delta_1, \delta_2, \delta_3, e$ shown in Figure 1. It follows that the corresponding V-arithmetic chamber $\mathcal{M}$ satisfies the following condition: the 2-dimensional face $\mathcal{M}_e$ of $\mathcal{M}$ which is perpendicular to e is an open fundamental triangle $\mathcal{M}_2$ bounded by three lines perpendicular to

$$P(\mathcal{M}_2) = \{\delta_1, \delta_2, \widetilde{\delta}_3\}$$



Figure 1: 3-dimensional graph $\Gamma_6^{(4)}$

for

$$\widetilde{\delta}_3 = \frac{\delta_3 + \cos(\pi/m)e}{\sin(\pi/m)}$$

(see Figure 1). It has one angle π/k, $k \geq 3$, defined by δ_1, δ_2. All its other sides don't intersect. Planes $\mathcal{H}_{\delta_1}$ and $\mathcal{H}_{\delta_2}$ are perpendicular to $\mathcal{H}_e$, and $\mathcal{H}_{\delta_3}$ has angle π/m, $m \geq 3$, with the plane $\mathcal{H}_e$.

Definition 2.4. *We denote by* $\Gamma_6^{(4)}(14)$ *the set of all V-arithmetic 3-dimensional graphs* $\Gamma_6^{(4)}$ *(or the corresponding 3-dimensional V-arithmetic chambers) of minimality* 14*. Thus, inequalities* $2 < a_{ij} < 14$ *satisfy. We denote by* $\mathcal{F}\Gamma_6^{(4)}(14)$ *the set of all their ground fields.*

Let us consider V-arithmetic 3-dimensional chambers which are defined by the Gram graphs $\Gamma_1^{(5)}$ with 5 vertices $\delta_1, \delta_2, \delta_3, \delta_4, e$ shown in Figure 2. It follows that the corresponding V-arithmetic chamber $\mathcal{M}$ satisfies the following condition: the 2-dimensional face $\mathcal{M}_e$ of $\mathcal{M}$ which is perpendicular to e is an open fundamental quadrangle $\mathcal{M}_2$ bounded by four lines perpendicular to

$$P(\mathcal{M}_2) = \{\widetilde{\delta}_1, \delta_2, \widetilde{\delta}_3, \delta_4\}$$

for

$$\widetilde{\delta}_1 = \frac{\delta_1 + \cos(\pi/m_1)e}{\sin(\pi/m_1)}, \quad \widetilde{\delta}_3 = \frac{\delta_3 + \cos(\pi/m_3)e}{\sin(\pi/m_3)}$$

(see Figure 2). It has two right angles defined by $\widetilde{\delta}_1, \delta_2$ and $\widetilde{\delta}_3, \delta_4$. All its other sides don't intersect. Planes $\mathcal{H}_{\delta_2}$ and $\mathcal{H}_{\delta_4}$ are perpendicular to $\mathcal{H}_e$, the planes $\mathcal{H}_{\delta_1}$ and $\mathcal{H}_{\delta_3}$ have angles π/m_1, $m_1 \geq 3$, and π/m_3, $m_3 \geq 3$, with the plane $\mathcal{H}_e$ respectively.



Figure 2: 3-dimensional graph $\Gamma_1^{(5)}$

Definition 2.5. *We denote by* $\Gamma_1^{(5)}(14)$ *the set of all V-arithmetic 3-dimensional graphs* $\Gamma_1^{(5)}$ *(or the corresponding 3-dimensional V-arithmetic chambers) of minimality* 14*. Thus, inequalities* $2 < a_{ij} < 14$ *satisfy. We denote by* $\mathcal{F}\Gamma_1^{(5)}(14)$ *the set of all their ground fields.*

Let us consider V-arithmetic 3-dimensional chambers which are defined by the Gram graphs $\Gamma_4^{(6)}$ with 6 vertices $\delta_1, \delta_2, \delta_3, \delta_4, \delta_5, e$ shown in Figure 3. It follows that the corresponding V-arithmetic chamber $\mathcal{M}$ satisfies the following condition: the 2-dimensional face $\mathcal{M}_e$ of $\mathcal{M}$ which is perpendicular to e is an open fundamental pentagon $\mathcal{M}_2$ bounded by 5 lines perpendicular to

$$P(\mathcal{M}_2) = \{\delta_1, \delta_2, \delta_3, \widetilde{\delta}_4, \delta_5\}$$

for

$$\widetilde{\delta}_4 = \frac{\delta_1 + \cos(\pi/m)e}{\sin(\pi/m)}$$

(see Figure 3). It has four consecutive right angles defined by δ_2, δ_3, $\widetilde{\delta}_4$, δ_5, δ_1 respectively. Its two consecutive sides perpendicular to δ_1 and δ_2 don't intersect. All planes $\mathcal{H}_{\delta_i}$ are perpendicular to $\mathcal{H}_e$ except the plane $\mathcal{H}_{\delta_4}$ which has the angle π/m with the plane $\mathcal{H}_e$.

Definition 2.6. *We denote by* $\Gamma_4^{(6)}(14)$ *the set of all V-arithmetic 3-dimensional graphs* $\Gamma_4^{(6)}$ *(or the corresponding 3-dimensional V-arithmetic chambers) of minimality* 14*. Thus, inequalities* $2 < a_{ij} < 14$ *satisfy. We denote by* $\mathcal{F}\Gamma_4^{(6)}(14)$ *the set of all their ground fields.*

We have

Theorem 2.7. *The sets of V-arithmetic graphs* $\Gamma_6^{(4)}(14)$*,* $\Gamma_1^{(5)}(14)$ *and* $\Gamma_4^{(6)}(14)$ *are finite.*



Figure 3: 3-dimensional graph $\Gamma_4^{(6)}$

Degree of any field from $\mathcal{F}\Gamma_6^{(4)}(14)$ *is bounded by* 56.
Degree of any field from $\mathcal{F}\Gamma_1^{(5)}(14)$ *is bounded by* 909.
Degree of any field from $\mathcal{F}\Gamma_4^{(6)}(14)$ *is bounded by* 99.

Proof. The proof requires long considerations and calculations. It will be given in a special Section 3. ☐

We have the following main result of the paper.

Theorem 2.8. *In dimensions* $n \geq 3$*, the ground field of any arithmetic hyperbolic reflection group belongs to one of finite sets of fields* $\mathcal{F}L^4$*,* $\mathcal{F}T$*,* $\mathcal{F}\Gamma_i^{(4)}(14)$*,* $1 \leq i \leq 5$*,* $\mathcal{F}\Gamma_{2,4}(14)$ *(fields in dimension* ≥ 6*), and* $\mathcal{F}\Gamma_1^{(6)}(14)$*,* $\mathcal{F}\Gamma_2^{(6)}(14)$*,* $\mathcal{F}\Gamma_3^{(6)}(14)$*,* $\mathcal{F}\Gamma_1^{(7)}(14)$*,* $\mathcal{F}\Gamma_2^{(7)}(14)$*,* $\mathcal{F}\Gamma_{2,5}(14)$ *(additional fields for the dimensions* 4 *and* 5*), and* $\mathcal{F}\Gamma_6^{(4)}(14)$*,* $\mathcal{F}\Gamma_1^{(5)}(14)$*,* $\mathcal{F}\Gamma_4^{(6)}(14)$*,* $\mathcal{F}\Gamma_2(14)$ *(additional fields for the dimension* 3*).*

In particular, its degree is bounded by 909.

Proof. By [20], if $n \geq 4$, the ground field $\mathbb{K}$ belongs to one of sets $\mathcal{F}L^4$, $\mathcal{F}T$, $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 5$, $\mathcal{F}\Gamma_{2,4}(14)$, $\mathcal{F}\Gamma_1^{(6)}(14)$, $\mathcal{F}\Gamma_2^{(6)}(14)$, $\mathcal{F}\Gamma_3^{(6)}(14)$, $\mathcal{F}\Gamma_1^{(7)}(14)$, $\mathcal{F}\Gamma_2^{(7)}(14)$, $\mathcal{F}\Gamma_{2,5}(14)$. Thus, further we can assume that the ground field $\mathbb{K}$ does not belong to these sets, and the dimension is equal to $n = 3$.

Let W be an arithmetic hyperbolic reflection group of dimension $n = 3$, $\mathcal{M}$ is its fundamental chamber, and $P(\mathcal{M})$ is the set of all vectors with square -2 which are perpendicular to codimension one faces of $\mathcal{M}$ and directed outward. For $\delta \in P(\mathcal{M})$ we denote by $\mathcal{H}_\delta$ and $\mathcal{M}_\delta$ the hyperplane and the codimension one face $\mathcal{M} \cap \mathcal{H}_\delta$ respectively which is perpendicular to δ.

By [14], there exists $e \in P(\mathcal{M})$ which defines a narrow face $\mathcal{M}_e$ of $\mathcal{M}$ of minimality 14. It means that $\delta_1 \cdot \delta_2 < 14$ for any $\delta_1, \delta_2 \in P(\mathcal{M}, e) \subset P(\mathcal{M})$. Here

$$P(\mathcal{M}, e) = \{\delta \in P(\mathcal{M}) \mid \mathcal{H}_\delta \cap \mathcal{H}_e \neq \emptyset\}.$$

Let us assume that $e \cdot \delta = 0$ for any $\delta \in P(\mathcal{M}, e) - \{e\}$, equivalently, all neighbouring 2-dimensional faces of $\mathcal{M}$ to the polygon $\mathcal{M}_e$ are perpendicular to $\mathcal{M}_e$. Then $\mathcal{M}_e$ has $P(\mathcal{M}_e) = P(\mathcal{M}, e) - \{e\}$, and $\mathcal{M}_e$ is the fundamental polygon for arithmetic hyperbolic plane reflection group with the same ground field $\mathbb{K}$ as for W. Since $P(\mathcal{M}_e) \subset P(\mathcal{M})$, it has minimality 14. Then $\mathbb{K} \in \mathcal{F}\Gamma_2(14)$ as required.

Thus, further we assume that $e \cdot \delta = 2\cos(\pi/m) > 0$ for one of $\delta \in P(\mathcal{M}, e) - \{e\}$, equivalently, one of neighbouring to the polygon $\mathcal{M}_e$ two-dimensional faces $\mathcal{M}_\delta$ of $\mathcal{M}$ is not perpendicular to $\mathcal{M}_e$.

All cases when the polygon $\mathcal{M}_e$ has less than 6 sides were considered in [19] and [20]. It was shown that then the ground field $\mathbb{K}$ belongs to one of sets of fields $\mathcal{F}L^4$, $\mathcal{F}T$, $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 5$, $\mathcal{F}\Gamma_{2,4}(14)$, $\mathcal{F}\Gamma_1^{(6)}(14)$, $\mathcal{F}\Gamma_2^{(6)}(14)$, $\mathcal{F}\Gamma_3^{(6)}(14)$, $\mathcal{F}\Gamma_1^{(7)}(14)$, $\mathcal{F}\Gamma_2^{(7)}(14)$, $\mathcal{F}\Gamma_{2,5}(14)$. Thus, further we additionally assume that $\mathcal{M}_e$ has at least 6 sides and the ground field $\mathbb{K}$ does not belong to any of these sets of fields.

By [19, Lemma 4.3], if $\mathbb{K}$ does not belong to $\mathcal{F}L^4$, $\mathcal{F}T$ and $\mathcal{F}\Gamma_i^{(4)}(14)$, $1 \leq i \leq 4$, then the Coxeter graph $C(v)$ of any vertex $v \in \mathcal{M}_e$ has all connected components having only one or two vertices. If additionally $\mathbb{K}$ does not belong to $\mathcal{F}\Gamma_5^{(4)}(14)$, then the hyperbolic connected component of the edge graph $\Gamma(r)$ defined by any edge $r = v_1v_2 \subset \mathcal{M}_e$ has ≤ 3 vertices. Further we mark these facts as (*).

By (*), both angles of $\mathcal{M}_e$ at the edge $\mathcal{M}_{e,\delta}$ perpendicular to δ are right. Moreover, if $f_1, f_2 \in P(\mathcal{M}, e)$ define two neighbouring edges of the edge $\mathcal{M}_{e,\delta}$ of $\mathcal{M}_e$, then $e \cdot f_1 = e \cdot f_2 = \delta \cdot f_1 = \delta \cdot f_2 = 0$.

Assume that the polygon $\mathcal{M}_e$ has a non-right angle with edges perpendicular to $\delta_1, \delta_2 \in P(\mathcal{M}, e)$. By (*), then $\delta_1 \cdot \delta_2 = 2\cos(\pi/k) > 0$, and $\{\delta_1, \delta_2\}$ are perpendicular to $\{e, \delta\}$. Then $\delta_1, \delta_2, \delta_3 = \delta$ and e have Gram graph $\Gamma_6^{(4)}(14)$, and the ground $\mathbb{K}$ belongs to $\mathcal{F}\Gamma_6^{(4)}(14)$ as required.

Now assume that all angles of the polygon $\mathcal{M}_e$ are right and there exist two elements $\delta_1, \delta_3 \in P(\mathcal{M}, e) - \{e\}$ such that $e \cdot \delta_1 = 2\cos(\pi/m_1)$ and $e \cdot \delta_3 = 2\cos(\pi/m_3)$. By (*), then $\delta_1 \cdot \delta_3 > 0$, and δ_1, δ_3 are perpendicular to two not consecutive edges of the polygon $\mathcal{M}_e$. Since $\mathcal{M}_e$ has more than 5 vertices, we can find two their neighbouring edges perpendicular to $\delta_2, \delta_4 \in P(\mathcal{M}, e)$ such that the Gram graph of $\delta_1, \delta_2, \delta_3, \delta_4, e$ is $\Gamma_1^{(5)}$. Then the ground field belongs to $\mathcal{F}\Gamma_1^{(5)}(14)$ as required.

Now assume that all angles of the polygon $\mathcal{M}_e$ are right and there exists only one element $\delta_4 \in P(\mathcal{M}, e) - \{e\}$ such that $\delta_4 \cdot e \neq 0$. Then $\delta_4 \cdot e = 2\cos(\pi/m) > 0$. Since $\mathcal{M}_e$ has at least 6 sides, we can find 5 consecutive sides of $\mathcal{M}_e$ perpendicular to $\delta_1, \delta_5, \delta_4, \delta_3, \delta_2 \in P(\mathcal{M}, e)$ such that Gram graph of $\delta_1, \delta_2, \delta_3, \delta_4, \delta_5, e$ is $\Gamma_4^{(6)}$. Then the ground field $\mathbb{K}$ belongs to $\mathcal{F}\Gamma_4^{(6)}(14)$ as required.

This finishes the proof of the theorem. $\square$

3 V-arithmetic 3-graphs $\Gamma_6^{(4)}(14)$, $\Gamma_1^{(5)}(14)$, $\Gamma_4^{(6)}(14)$ and their fields

Here we prove Theorem 2.7.

3.1 Some general results.

We use the following general results from [15].

Theorem 3.1. *([15, Theorem 1.2.1]) Let $\mathbb{F}$ be a totally real algebraic number field, and let each embedding $\sigma : \mathbb{F} \to \mathbb{R}$ corresponds to an interval $[a_\sigma, b_\sigma]$ in $\mathbb{R}$ where*

$$\prod_\sigma \frac{b_\sigma - a_\sigma}{4} < 1.$$

In addition, let the natural number m and the intervals $[s_1, t_1], \dots, [s_m, t_m]$ in $\mathbb{R}$ be fixed. Then there exists a constant $N(s_i, t_i)$ such that, if α is a totally real algebraic integer and if the following inequalities hold for the embeddings $\tau : \mathbb{F}(\alpha) \to \mathbb{R}$:

$$s_i \le \tau(\alpha) \le t_i \quad for \quad \tau = \tau_1, \dots, \tau_m,$$

$$a_{\tau|\mathbb{F}} \le \tau(\alpha) \le b_{\tau|\mathbb{F}} \quad for \quad \tau \ne \tau_1, \dots, \tau_m,$$

then

$$[\mathbb{F}(\alpha) : \mathbb{F}] \le N(s_i, t_i).$$

Theorem 3.2. *([15, Theorem 1.2.2]) Under the conditions of Theorem 3.1, $N(s_i, t_i)$ can be taken to be $N(s_i, t_i) = N_0$, where N_0 is the least natural number solution of the inequality*

$$N_0 M \ln(1/R) - M \ln(N_0 + 1) - \ln B \ge \ln S. \tag{1}$$

Here

$$M = [\mathbb{F} : \mathbb{Q}], \quad B = 2\sqrt{|\mathrm{discr}\ \mathbb{F}|}; \tag{2}$$

$$R = \sqrt{\prod_\sigma \frac{b_\sigma - a_\sigma}{4}}, \quad S = \prod_{i=1}^m \frac{2er_i}{b_{\sigma_i} - a_{\sigma_i}} \tag{3}$$

where

$$\sigma_i = \tau_i|\mathbb{F}, \quad r_i = \max\{|b_i - a_{\sigma_i}|, |b_{\sigma_i} - a_i|\}. \tag{4}$$

We note that the proof of Theorems 3.1 and 3.2 uses a variant of Fekete's Theorem (1923) about existence of non-zero integer polynomials of bounded degree which differ only slightly from zero on appropriate intervals. See [15, Theorem 1.1.1].

Below we will apply these results in two cases which are very similar to used in [19, Sec. 5.5] and [15]. Cases 1 and 2 below are natural generalizations of Cases 1 and 2 which were considered in [20, Sec. 3.1].

Case 1. For a natural $l \geq 3$ we denote $\mathbb{F}_l = \mathbb{Q}(\cos(2\pi/l))$. We consider a totally real algebraic number field $\mathbb{K}$ where $\mathbb{F}_l \subset \mathbb{K} = \mathbb{Q}(\alpha)$, and the algebraic integer α satisfies

$$-a_1\sigma(\sin^2(\pi/l)) < \sigma(\alpha) < a_2\sigma(\sin^2(\pi/l)) \tag{5}$$

for all $\sigma : \mathbb{K} \to \mathbb{R}$ such that $\sigma \neq \sigma^{(+)}$, and

$$b_1 < \sigma^{(+)}(\alpha) < b_2 \tag{6}$$

where $\sigma^{(+)} : \mathbb{K} \to \mathbb{R}$ is the identity. We assume that $a_1 \geq 0$, $a_2 \geq 0$ and $0 < a = \max\{a_1, a_2\} < 4$. We assume that $b_1 < b_2$ and denote $b = \max\{|b_1|, |b_2|\}$. Also we assume that $a \leq b$. We want to estimate $[\mathbb{K} : \mathbb{F}_l] = N_0$ and $N = [\mathbb{K} : \mathbb{Q}] = N_0 \cdot [\mathbb{F}_l : \mathbb{Q}]$ from above.

For $l \geq 3$, we have $[\mathbb{F}_l : \mathbb{Q}] = \varphi(l)/2$ where $\varphi(l)$ is the Euler function, and $N_{\mathbb{F}_l/\mathbb{Q}}(\sin^2(\pi/l)) = \gamma(l)/4^{\varphi(l)/2}$ where

$$N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2(\pi/l)) = \gamma(l) = \begin{cases} p & \text{if } l = p^t > 2 \text{ where } p \text{ is prime,} \\ 1 & \text{otherwise.} \end{cases} \tag{7}$$

We have

$$\frac{ba^N |N_{\mathbb{K}/\mathbb{Q}}(\sin^2(\pi/l))|}{a\sin^2(\pi/l)} > |N_{\mathbb{K}/\mathbb{Q}}(\alpha)| \geq 1$$

and

$$\frac{b(a/4)^N \gamma(l)^{2N/\varphi(l)}}{a\sin^2(\pi/l)} > 1.$$

Equivalently, we have

$$N\left(\ln\frac{2}{\sqrt{a}} - \frac{\ln\gamma(l)}{\varphi(l)}\right) < \ln\sqrt{\frac{b}{a}} - \ln\sin\frac{\pi}{l} \ \text{ and } \ (\varphi(l)/2)|N. \tag{8}$$

Since $\gamma(l) \leq l$, $\varphi(l) \geq Cl/\ln(\ln l)$ for $l \geq 6$ where $C = \varphi(6)\ln(\ln 6)/6 \geq 0.194399$, $\sin(\pi/l) \leq \pi/l$ for $l \geq 3$, there exists only finite number of $l \geq 3$ such that (8) has solutions $N \in \mathbb{N}$.

More exactly, there exists only finite number of *exceptional* $l \geq 3$ such that

$$\ln\frac{2}{\sqrt{a}} - \frac{\ln\gamma(l)}{\varphi(l)} \leq 0. \tag{9}$$

All non-exceptional l satisfy the inequality

$$(\varphi(l)/2)\left(\ln\frac{2}{\sqrt{a}}-\frac{\ln\gamma(l)}{\varphi(l)}\right)<\ln\sqrt{\frac{b}{a}}-\ln\sin\frac{\pi}{l}. \tag{10}$$

Remark that exceptional l also satisfy this inequality.

If $\gamma(l)=1$, (10) implies that l satisfies the inequality

$$(C/2)\ln(2/\sqrt{a})\,l<\Big(\ln l+\ln(\sqrt{(b/a)}/\pi)\Big)\ln\ln l\,. \tag{11}$$

It follows that

$$l<L_0 \tag{12}$$

where $L_0>3$ satisfies

$$(C/2)\ln(2/\sqrt{a})\,L_0\geq\Big(\ln L_0+\ln(\sqrt{(b/a)}/\pi)\Big)\ln\ln L_0\,. \tag{13}$$

If $l=p^t$ where p is prime, (10) implies that l satisfies the inequality

$$(C/2)\Delta(a)\,l<\Big(\ln l+\ln(\sqrt{(b/a)}/\pi)\Big)\ln\ln l \tag{14}$$

where

$$\Delta(a)=\min_{l=p^t\geq L_0}\left\{\ln\frac{2}{\sqrt{a}}-\frac{\ln\gamma(l)}{\varphi(l)}>0\right\}. \tag{15}$$

It follows that

$$l<L_1 \tag{16}$$

where $L_1\geq L_0$ is a solution of the inequality

$$(C/2)\Delta(a)\,L_1\geq\Big(\ln L_1+\ln(\sqrt{(b/a)}/\pi)\Big)\ln\ln L_1\,. \tag{17}$$

Thus, to find all non-exceptional l satisfying (10), we should check (10) for all l such that $3\leq l<L_1$, moreover, if $L_0\leq l<L_1$, we can assume that $l=p^t$. Their number is finite, and all of them can be effectively found.

For non-exceptional l satisfying (10), we obtain bounds

$$N_0=[\mathbb{K}:\mathbb{F}_l]\leq\left[\frac{\ln\sqrt{b/a}-\ln\sin(\pi/l)}{(\varphi(l)/2)\,(\ln(2/\sqrt{a})-(\ln\gamma(l))/\varphi(l))}\right] \tag{18}$$

and

$$N=[\mathbb{K}:\mathbb{Q}]\leq\left[\frac{\ln\sqrt{b/a}-\ln\sin(\pi/l)}{(\varphi(l)/2)\,(\ln(2/\sqrt{a})-(\ln\gamma(l))/\varphi(l))}\right]\cdot(\varphi(l)/2)\,. \tag{19}$$

This using of the norm, we call the *Method B* (like in [19, Sec. 5.5]).

On the other hand, for fixed l, we obtain a bound for N_0 using Theorems 3.1 and 3.2 applied to $\mathbb{F} = \mathbb{F}_l$ and α. We can take

$$R = \sqrt{|N_{\mathbb{F}_l/\mathbb{Q}}(\sin^2(\pi/l))|\,((a_1+a_2)/4)^{\varphi(l)/2}} = \left(\frac{\gamma(l)^{1/\varphi(l)}(a_1+a_2)^{1/2}}{4}\right)^{\varphi(l)/2}, \tag{20}$$

where

$$R < 1 \ \text{ if and only if } \ \ln\frac{4}{\sqrt{a_1+a_2}} - \frac{\ln\gamma(l)}{\varphi(l)} > 0\,, \tag{21}$$

$$M = [\mathbb{F}_l : \mathbb{Q}] = \frac{\varphi(l)}{2}, \quad B = 2\sqrt{|\operatorname{discr}\mathbb{F}_l|} \tag{22}$$

where the discriminant $|\operatorname{discr}\mathbb{F}_l|$ is given in (70), and

$$S = \frac{2e\max\{a_2, b_2, a_2-b_1, a_1, -b_1, b_2+a_1\}}{(a_1+a_2)\sin^2(\pi/l)}. \tag{23}$$

Then $[\mathbb{K} : \mathbb{F}_l] \le n_0$ and $[\mathbb{K} : \mathbb{Q}] \le n_0\varphi(l)/2$ where n_0 is the least natural solution of the inequality (1)

$$n_0 M\ln(1/R) - M\ln(n_0+1) - \ln B \ge \ln S. \tag{24}$$

In particular, this gives a bound for $[\mathbb{K} : \mathbb{Q}]$ for exceptional l satisfying (21) and improves the bound (18) for N_0 when it is poor, which also improves the bound for $[\mathbb{K} : \mathbb{Q}]$. This using of Theorems 3.1, 3.2, we call the *Method A* (like in [19, Sec. 5.5]).

We shall apply these Methods A and B to $\Gamma_4^{(6)}(14)$ in Sect. 3.4.

Case 2. For natural $k \ge s \ge 3$, we denote $\mathbb{F}_{k,s} = \mathbb{Q}(\cos(2\pi/k), \cos(2\pi/s))$. We consider a totally real algebraic number field $\mathbb{K}$ where $\mathbb{F}_{k,s} \subset \mathbb{K} = \mathbb{Q}(\alpha)$, and the algebraic integer α satisfies

$$-a_1\sigma(\sin^2(\pi/k)\sin^2(\pi/s)) < \sigma(\alpha) < a_2\sigma(\sin^2(\pi/k)\sin^2(\pi/s)) \tag{25}$$

for all $\sigma : \mathbb{K} \to \mathbb{R}$ such that $\sigma \ne \sigma^{(+)}$, and

$$b_1 < \sigma^{(+)}(\alpha) < b_2 \tag{26}$$

where $\sigma^{(+)} : \mathbb{K} \to \mathbb{R}$ is the identity. We assume that $a_1 \ge 0$, $a_2 \ge 0$ and $0 < a = \max\{a_1, a_2\} < 16$. We assume that $b_1 < b_2$ and denote $b = \max\{|b_1|, |b_2|\}$. Also we assume that $a \le b$. We want to estimate $[\mathbb{K} : \mathbb{F}_{k,s}] = N_0$ and $N = [\mathbb{K} : \mathbb{Q}] = N_0[\mathbb{F}_{k,s} : \mathbb{Q}]$ for non-exceptional k and s where $l \ge 3$ *is called exceptional* if

$$\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(l)}{\varphi(l)} \le 0. \tag{27}$$

We also assume that $k \ge s \ge s_0 \ge 3$ where $s_0 \ge 3$ is fixed.

We have $[\mathbb{F}_{k,s}:\mathbb{Q}] = \varphi([k,s])/2\rho(k,s)$ where $\rho(k,s)=1$ or 2 is given in (71), and $N_{\mathbb{F}_l/\mathbb{Q}}(\sin^2(\pi/l)) = \gamma(l)/4^{\varphi(l)/2}$ where $\gamma(l)$ is given in (7). We have

$$\frac{ba^N|N_{\mathbb{K}/\mathbb{Q}}(\sin^2(\pi/k))||N_{\mathbb{K}/\mathbb{Q}}(\sin^2(\pi/s))|}{a\sin^2(\pi/k)\sin^2(\pi/s)} > |N_{\mathbb{K}/\mathbb{Q}}(\alpha)| \geq 1$$

and

$$\frac{b(a/16)^N\gamma(k)^{2N/\varphi(k)}\gamma(s)^{2N/\varphi(s)}}{a\sin^2(\pi/k)\sin^2(\pi/s)} > 1.$$

Equivalently, we obtain

$$N\left(\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(k)}{\varphi(k)} - \frac{\ln\gamma(s)}{\varphi(s)}\right) < \ln\sqrt{\frac{b}{a}} - \ln\sin\frac{\pi}{k} - \ln\sin\frac{\pi}{s} \text{ and } \frac{\varphi([k,s])}{2\rho(k,s)} \mid N. \tag{28}$$

Since $\gamma(l) \leq l$, $\varphi(l) \geq Cl/\ln(\ln l)$ for $l \geq 6$ where $C = \varphi(6)\ln(\ln 6)/6$, $\sin(\pi/l) \leq \pi/l$ for $l \geq 3$, there exists only finite number of pairs (k,s) such that (28) has solutions $N \in \mathbb{N}$ where both k and s are non-exceptional.

More exactly, there exists only finite number of *exceptional pairs* (k,s) where a pair (k,s) (consisting of non-exceptional k and s) is called exceptional if

$$\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(k)}{\varphi(k)} - \frac{\ln\gamma(s)}{\varphi(s)} \leq 0. \tag{29}$$

All non-exceptional pairs (k,s) satisfying (28) satisfy the inequality

$$\frac{\varphi([k,s])}{2\rho(k,s)} \cdot \left(\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(k)}{\varphi(k)} - \frac{\ln\gamma(s)}{\varphi(s)}\right) < \ln\sqrt{\frac{b}{a}} - \ln\sin\frac{\pi}{k} - \ln\sin\frac{\pi}{s}\ . \tag{30}$$

Remark that exceptional pairs (k,s) also satisfy this inequality.

If $\gamma(k)=\gamma(s)=1$ and $k \geq s$, (30) implies

$$(C/2)\ln(4/\sqrt{a})k < \Big(2\ln k + \ln(\sqrt{(b/a)}/\pi^2)\Big)\ln\ln k. \tag{31}$$

It follows that

$$s_0 \leq s \leq k < K_0 \tag{32}$$

where $K_0 > 3$ satisfies

$$(C/2)\ln(4/\sqrt{a})K_0 \geq \Big(2\ln K_0 + \ln(\sqrt{(b/a)}/\pi^2)\Big)\ln\ln K_0. \tag{33}$$

If one of $\gamma(k)$, $\gamma(s)$ is not equal to 1, then (30) implies for non-exceptional pairs (k,s) that

$$(C/2)\Delta_1(a)k < \Big(2\ln k + \ln(\sqrt{(b/a)}/\pi^2)\Big)\ln\ln k \tag{34}$$

where

$$\Delta_1(a) = \min_{k\geq s\geq s_0, k\geq K_0}\left\{\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(s)}{\varphi(s)} - \frac{\ln\gamma(k)}{\varphi(k)}\ > 0\right\}. \tag{35}$$

It follows that

$$s_0 \le s \le k < K_1 \tag{36}$$

where $K_1 \ge K_0$ is a solution of the inequality

$$(C/2)\Delta_1(a)K_1 \ge \Big(2\ln K_1 + \ln\big(\sqrt{(b/a)}/\pi^2\big)\Big)\ln\ln K_1. \tag{37}$$

Thus, to find all non-exceptional pairs (k, s) satisfying (30), we should check (30) for all $s_0 \le s \le k < K_1$; moreover, if $K_0 \le k \le K_1$, we can assume that one of k and s is equal to p^t where p is prime. The number of such pairs is finite, and all of them can be effectively found.

For such non-exceptional pairs (k, s) satisfying (30), we obtain bounds

$$N_0 = [\mathbb{K} : \mathbb{F}_{k,s}] \le \left[\frac{\ln\sqrt{\frac{b}{a}} - \ln\sin\frac{\pi}{k} - \ln\sin\frac{\pi}{s}}{\frac{\varphi([k,s])}{2\rho(k,s)}\cdot\Big(\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(k)}{\varphi(k)} - \frac{\ln\gamma(s)}{\varphi(s)}\Big)}\right] \tag{38}$$

and

$$N = [\mathbb{K} : \mathbb{Q}] \le \left[\frac{\ln\sqrt{\frac{b}{a}} - \ln\sin\frac{\pi}{k} - \ln\sin\frac{\pi}{s}}{\frac{\varphi([k,s])}{2\rho(k,s)}\cdot\Big(\ln\frac{4}{\sqrt{a}} - \frac{\ln\gamma(k)}{\varphi(k)} - \frac{\ln\gamma(s)}{\varphi(s)}\Big)}\right]\cdot\frac{\varphi([k,s])}{2\rho(k,s)}\,. \tag{39}$$

This using of the norm, we call the *Method B* (like in [19, Sec. 5.5]).

On the other hand, for a fixed pair (k, s), we can obtain a bound for N_0 using Theorems 3.1 and 3.2 applied to $\mathbb{F} = \mathbb{F}_{k,s}$ and α. We can take

$$R = \sqrt{|N_{\mathbb{F}_{k,s}/\mathbb{Q}}(\sin^2\frac{\pi}{k}\sin^2\frac{\pi}{s})|\left(\frac{a_1+a_2}{4}\right)^{\frac{\varphi([k,s])}{2\rho(k,s)}}} =$$

$$= \left(\frac{\gamma(k)^{\frac{1}{\varphi(k)}}\gamma(s)^{\frac{1}{\varphi(s)}}(a_1+a_2)^{\frac{1}{2}}}{8}\right)^{\frac{\varphi([k,s])}{2\rho(k,s)}} \tag{40}$$

where

$$R < 1 \text{ if and only if } \ln\frac{8}{\sqrt{a_1+a_2}} - \frac{\ln\gamma(k)}{\varphi(k)} - \frac{\ln\gamma(s)}{\varphi(s)} > 0, \tag{41}$$

$$M = [\mathbb{F}_{k,s} : \mathbb{Q}] = \frac{\varphi([k,s])}{2\rho(k,s)},\quad B = 2\sqrt{|\mathrm{discr}\,\mathbb{F}_{k,s}|} \tag{42}$$

where the discriminant $|\mathrm{discr}\,\mathbb{F}_{k,s}|$ is given in (73) and (74), and

$$S = \frac{2e\max\{a_2, b_2, a_2 - b_1, a_1, -b_1, b_2 + a_1\}}{(a_1+a_2)\sin^2(\pi/s)\sin^2(\pi/k)}. \tag{43}$$

For all pairs (k, s) satisfying (41), we obtain the bounds $[\mathbb{K} : \mathbb{F}_{k,s}] \le n_0$ and $[\mathbb{K} : \mathbb{Q}] \le n_0\varphi([k, s])/(2\rho(k, s))$ where n_0 is the least natural solution of the inequality (1),

$$n_0 M\ln(1/R) - M\ln(n_0+1) - \ln B \ge \ln S.$$



Figure 4: The graph of $\mathcal{M}_2$ for $\Gamma_1^{(5)}$

For $a < 16$ and $k, s \geq 3$, all pairs (k, s), except finite number, satisfy (41), and we can apply this method to all these pairs. In particular, this gives a bound for $[\mathbb{K} : \mathbb{Q}]$ for all exceptional pairs (k, s) satisfying (41), and it improves the bound (38) for N_0 when it is poor, which also improves the bound for $[\mathbb{K} : \mathbb{Q}]$. This using of Theorems 3.1, 3.2, we call the *Method A* (like in [19, Sec. 5.5]).

We apply these Methods A and B to $\Gamma_6^{(4)}(14)$ in Sect. 3.2 and to $\Gamma_1^{(5)}(14)$ in Sect. 3.3.

3.2 V-arithmetic 3-graphs $\Gamma_6^{(4)}(14)$ and their fields.

Here we consider V-arithmetic 3-dimensional graphs $\Gamma_6^{(4)}(14)$ (see Figure 1) and their fields.

This case had been considered in [20, Sec. 3.1] where the bound for degrees of fields $\mathbb{K}$ from $\mathcal{F}\Gamma_1^{(6)}(14)$ was obtained. To get this bound, in [20, Sec. 3.2] the bound for fields defined by the subgraph $\Gamma_6^{(4)}(14)$ of this graph was obtained. This uses methods A and B of Case 2 in Sect. 3.1 applied to $a_1 = 0$, $a_2 = 4$, $b_1 = 12$ and $b_2 = 28^2$. The bound is $[\mathbb{K} : \mathbb{Q}] \leq 56$.

3.3 V-arithmetic 3-graphs $\Gamma_1^{(5)}(14)$ and their fields.

Here we consider V-arithmetic 3-dimensional graphs $\Gamma_1^{(5)}(14)$ (see Figure 2) and their fields.

First, let us consider the corresponding plane graph defined by $\beta_1 = \widetilde{\delta_1}$, $\beta_2 = \delta_2$, $\beta_3 = \widetilde{\delta_3}$, $\beta_4 = \delta_4$ which give $P(\mathcal{M}_2)$. We denote $b_{ij} = \beta_i \cdot \beta_j$ when it is not 0. This graph is given in Figure 4.

Any three elements from $\beta_1, \ldots, \beta_4$ generate the form defining the hyperbolic plane. Thus the determinant of their Gram matrix must be positive for geometric embedding $\sigma^{(+)}$ and must be negative for $\sigma \neq \sigma^{(+)}$. For example, for $\beta_1, \beta_2, \beta_3$ it is equal to $-8 + 2b_{13}^2 + 2b_{23}^2$. Thus, for σ we obtain inequalities

$b_{13}^2+b_{23}^2<4$. Moreover, the determinant

$$16+b_{13}^2b_{24}^2+b_{14}^2b_{23}^2-4b_{13}^2-4b_{14}^2-4b_{23}^2-4b_{24}^2-2b_{13}b_{14}b_{23}b_{24}$$

of the Gram matrix of all four elements $\beta_1,\dots,\beta_4$ is 0. Combining all these conditions, we obtain the following conditions on $\mathcal{M}_2$ for $\sigma\neq\sigma^{(+)}$:

$$\begin{cases} b_{13}b_{14}b_{23}b_{24}=8-2b_{13}^2-2b_{14}^2-2b_{23}^2-2b_{24}^2+(b_{13}^2b_{24}^2+b_{14}^2b_{23}^2)/2\\ b_{13}^2+b_{23}^2<4\\ b_{23}^2+b_{24}^2<4\\ b_{24}^2+b_{14}^2<4\\ b_{14}^2+b_{13}^2<4\ . \end{cases} \tag{44}$$

It is easy to find minimum and maximum of $b_{13}b_{14}b_{23}b_{24}$ under the closure of these conditions which shows that

$$-4<\sigma(b_{13}b_{14}b_{23}b_{24})\leq 1. \tag{45}$$

Here minimum is achieved for $b_{ij}=\pm\sqrt{2}$ where the number of $(-)$ is odd, and maximum is achieved for $b_{ij}=\pm1$ where the number of $(-)$ is even. From expressions of β_i using δ_i and e, we get

$$b_{13}b_{14}b_{23}b_{24}=\frac{a_{13}a_{14}a_{23}a_{24}+2\cos\frac{\pi}{m_1}\cos\frac{\pi}{m_3}a_{14}a_{23}a_{24}}{\sin^2\frac{\pi}{m_1}\sin^2\frac{\pi}{m_3}}. \tag{46}$$

We consider the algebraic integer $\alpha\in\mathbb{K}$ which is

$$\alpha=2a_{13}a_{14}a_{23}a_{24}+4\cos\frac{\pi}{m_1}\cos\frac{\pi}{m_3}a_{14}a_{23}a_{24}. \tag{47}$$

From (45) and (46), we get

$$-8\sigma(\sin^2\frac{\pi}{m_1}\sin^2\frac{\pi}{m_3})<\sigma(\alpha)\leq 2\sigma(\sin^2\frac{\pi}{m_1}\sin^2\frac{\pi}{m_3}). \tag{48}$$

For the geometric embedding $\sigma^{(+)}$, we have

$$2\cdot 2^4+2^3=5\cdot 2^3<\sigma^{(+)}(\alpha)<2\cdot 14^4+4\cdot 14^3=32\cdot 14^3. \tag{49}$$

It follows that $\mathbb{K}=\mathbb{Q}(\alpha)$. Since $8<16$, this case is similar to considered in [19, Sec. 5.5].

In this case, considering α from (47), we apply the methods A and B of Case 2 in Sec. 3.1 to $a_1=8$, $a_2=2$ (then $a=8$), $b_1=5\cdot 2^3$, $b_2=32\cdot 14^3$ (then $b=32\cdot 14^3$) and $k:=\max\{m_1,m_3\}$, $s:=\min\{m_1,m_3\}$ where $k\geq s\geq 3$, and $s_0=3$.

At first, we apply the Method B. Exceptional $l\geq 3$ satisfy (27) which is

$$\ln\sqrt{2}-\frac{\ln\gamma(l)}{\varphi(l)}\leq 0. \tag{50}$$

It follows that $l = 3, 4, 5$ are the only exceptional.

All exceptional pairs (k, s) where $k \geq s \geq 6$, that is when (29) which is

$$\ln \sqrt{2} - \frac{\ln \gamma(k)}{\varphi(k)} - \frac{\ln \gamma(s)}{\varphi(s)} \leq 0 \tag{51}$$

satisfies, are $(k, s = 7)$ where either $7 \leq k \leq 241$ is prime, or $k = 8$, 9, 16, 25, 27, 32, 49; $(k, s = 8)$ where $k = 8$, 9, 11, 13, 17; $(k, s = 9)$ where $k = 9$, 11, 13, 17, 19; $(k, s = 11)$ where $11 \leq k \leq 31$ is prime; $(k, s = 13)$ where $13 \leq k \leq 23$ is prime; $(k = 17, s = 17)$.

We can take $K_0 = 911$ in (33). Then (here we take $s_0 = 6$)

$$\Delta_1(8) = \ln \sqrt{2} - \frac{\ln 7}{6} - \frac{\ln 911}{910} \geq 0.0147667,$$

and $K_1 = 38563$ can be taken in (37). Checking (30) for $6 \leq s \leq k < 38563$, we obtain that $6 \leq s \leq 330$ and $6 \leq s \leq k \leq 5460$. Moreover, $s \leq k \leq 330$ for $20 \leq s \leq 330$. For all these pairs (k, s) satisfying (30) which is

$$\frac{\varphi([k, s])}{2\rho(k, s)} \cdot \left(\ln \sqrt{2} - \frac{\ln \gamma(k)}{\varphi(k)} - \frac{\ln \gamma(s)}{\varphi(s)} \right) < \ln \sqrt{4 \cdot 14^3} - \ln \sin \frac{\pi}{k} - \ln \sin \frac{\pi}{s}, \tag{52}$$

we obtain

$$[\mathbb{F}_{k,s} : \mathbb{Q}] = \frac{\varphi([k, s])}{2\rho(k, s)} \leq 909 \tag{53}$$

where 909 is achieved for $(k, s) = (607, 7)$. Moreover, for all these non-exceptional pairs (k, s) we obtain the bound (38) which is

$$N_0 = [\mathbb{K} : \mathbb{F}_{k,s}] \leq \left[\frac{\ln \sqrt{2} - \ln \sin \frac{\pi}{k} - \ln \sin \frac{\pi}{s}}{\frac{\varphi([k,s])}{2\rho(k,s)} \cdot \left(\ln \sqrt{4 \cdot 14^3} - \frac{\ln \gamma(k)}{\varphi(k)} - \frac{\ln \gamma(s)}{\varphi(s)} \right)} \right] \tag{54}$$

and finally we obtain the bound (39) which is

$$N = [\mathbb{K} : \mathbb{Q}] \leq \left[\frac{\ln \sqrt{2} - \ln \sin \frac{\pi}{k} - \ln \sin \frac{\pi}{s}}{\frac{\varphi([k,s])}{2\rho(k,s)} \cdot \left(\ln \sqrt{4 \cdot 14^3} - \frac{\ln \gamma(k)}{\varphi(k)} - \frac{\ln \gamma(s)}{\varphi(s)} \right)} \right] \cdot \frac{\varphi([k, s])}{2\rho(k, s)}\ . \tag{55}$$

If either a pair (k, s) is exceptional, or the right hand side of (55) is more than 909 (these are possible only for pairs (k, s) with $6 \leq s \leq 17$ and $s \leq k \leq 421$), we also apply to the pair (k, s) the method A of the Case 2 to improve the poor bound (54) of $N_0 = [\mathbb{K} : \mathbb{F}_{k,s}]$ for non-exceptional (k, s). We can apply this method to any pair (k, s) with $k \geq s \geq 6$ since (41) is valid if $a_1 + a_2 = 10$. We obtain that $[\mathbb{K} : \mathbb{Q}] \leq 909$ for all $k \geq s \geq 6$.

Let us assume that $s = 3, 4$ or 5 is exceptional. It means that either $m_1 = 3, 4, 5$ or $m_3 = 3, 4, 5$ for $\Gamma_1^{(5)}(14)$. For example, let $m_1 = 3, 4, 5$. Let us consider the V-arithmetic graph defined by e, δ_1 and δ_4. We denote $\alpha = a_{14}^2$ where the algebraic integer $a_{14} = \delta_1 \cdot \delta_4$. The determinant of the Gram matrix of e, δ_1



Figure 5: The graph of $\mathcal{M}_2$ for $\Gamma_4^{(6)}$

and δ_4 is equal to $2\alpha - 8\sin^2(\pi/m_1)$. It follows that $0 < \sigma(\alpha) < 4\sin^2(\pi/m_1)$ for $\sigma \neq \sigma^{(+)}$, and $4 < \sigma^{(+)}(\alpha) < 14^2$. Then $\mathbb{K} = \mathbb{Q}(\alpha)$ and $\mathbb{F}_{m_1} \subset \mathbb{K}$. Thus, we can apply the method A of Case 1 in Sec. 3.1 to $a_1 = 0$, $a_2 = 4$, $b_1 = 4$, $b_2 = 14^2$ and $l := m_1$ where $m_1 = 3, 4, 5$. We obtain that $[\mathbb{K} : \mathbb{Q}] \leq 76$ for $m_1 = 3$, $[\mathbb{K} : \mathbb{Q}] \leq 31$ for $m_1 = 4$, and $[\mathbb{K} : \mathbb{Q}] \leq 24$ for $m_1 = 5$.

Thus, finally, $[\mathbb{K} : \mathbb{Q}] \leq 909$ for all graphs $\Gamma_1^{(5)}(14)$.

3.4 V-arithmetic 3-graphs $\Gamma_4^{(6)}(14)$ and their ground fields

Here we consider V-arithmetic 3-dimensional graphs $\Gamma_4^{(6)}(14)$ and their fields.

First, let us consider the corresponding plane graph defined by $\beta_1 = \delta_1$, $\beta_2 = \delta_2$, $\beta_3 = \delta_3$, $\beta_4 = \widetilde{\delta}_4$, $\beta_5 = \delta_5$ which give $P(\mathcal{M}_2)$. We denote $b_{ij} = \beta_i \cdot \beta_j$ when it is not 0. We also denote $c = b_{12}$. One can consider it as kind of angle between the corresponding lines. This graph is given in Figure 5.

Considering determinants of Gram matrices of subsets of $\beta_1, \dots, \beta_5$, we obtain all equations of $\mathcal{M}_2$:

$$\begin{cases} 4b_{13}^2 = (4 - b_{14}^2)(4 - b_{35}^2) \\ 4b_{25}^2 = (4 - b_{24}^2)(4 - b_{35}^2) \\ 4b_{14}^2 + 4c^2 + 4cb_{14}b_{24} = (4 - b_{13}^2)(4 - b_{24}^2) \\ 4b_{24}^2 + 4c^2 + 4cb_{14}b_{24} = (4 - b_{14}^2)(4 - b_{25}^2) \\ b_{35}^2(4 - c^2) + 2cb_{13}b_{25}b_{35} + 4c^2 = (4 - b_{13}^2)(4 - b_{25}^2). \end{cases} \tag{56}$$

For $\sigma \neq \sigma^{(+)}$ they also satisfy inequalities: $b_{ij}^2 < 4$ for all b_{ij}, $c^2 < 4$. By direct calculation of minimum and maximum of $b_{13}b_{14}b_{24}b_{25}b_{35}$ for $0 \leq b_{ij}^2 \leq 4$ and $0 \leq c^2 \leq 4$ satisfying equations (56), we obtain that

$$-3.1 < \sigma(b_{13}b_{14}b_{24}b_{25}b_{35}) < 3.1\ . \tag{57}$$

Here minimum $-3.07\dots$ is achieved for $c = -0.39\dots$, $b_{14} = b_{24} = \pm 1.4\dots$, $b_{13} = \pm 1.166\dots$, $b_{25} = \pm 1.166\dots$, $b_{35} = \pm 1.1549\dots$ and $c = 0.39\dots$, $b_{14} =$

$-b_{24} = \pm 1.4\ldots$, $b_{13} = \pm 1.166\ldots$, $b_{25} = \pm 1.166\ldots$, $b_{35} = \pm 1.1549\ldots$. Here maximum $3.07\ldots$ is achieved for $c = -1.569\ldots$, $b_{14} = b_{24} = \pm 1.4\ldots$, $b_{13} = \pm 1.166\ldots$, $b_{25} = \pm 1.166\ldots$, $b_{35} = \pm 1.1549\ldots$ and $c = 1.569\ldots$, $b_{14} = -b_{24} = \pm 1.4\ldots$, $b_{13} = \pm 1.166\ldots$, $b_{25} = \pm 1.166\ldots$, $b_{35} = \pm 1.1549\ldots$.

From expressions of β_i using δ_i and e, we get

$$b_{13}b_{14}b_{24}b_{25}b_{35} = \frac{a_{13}a_{14}a_{24}a_{25}a_{35}}{\sin^2 \frac{\pi}{m}}. \tag{58}$$

We consider the algebraic integer $\alpha \in \mathbb{K}$ which is

$$\alpha = a_{13}a_{14}a_{24}a_{25}a_{35}. \tag{59}$$

From (57), we obtain

$$-3.1 \cdot \sigma(\sin^2 \frac{\pi}{m}) < \sigma(\alpha) < 3.1 \cdot \sigma(\sin^2 \frac{\pi}{m}). \tag{60}$$

For the geometric embedding $\sigma^{(+)}$ we have that

$$2^5 < \sigma^{(+)}(\alpha) < 14^5. \tag{61}$$

It follows that $\mathbb{K} = \mathbb{Q}(\alpha)$.

We can apply the methods A and B of Case 1 in Sec. 3.1 to $a_1 = 3.1$, $a_2 = 3.1$ (then $a = 3.1$), $b_1 = 2^5$, $b_2 = 14^5$ (then $b = 14^5$), and $l := m$.

At first, we apply the Method B. All exceptional $l \geq 3$ that is when (9) which is

$$\ln \frac{2}{\sqrt{3.1}} - \frac{\ln \gamma(l)}{\varphi(l)} \leq 0 \tag{62}$$

satisfies are $l = 3, 4, 5, 7, 8, 9, 11, 13, 17, 19, 23$.

We can take $L_0 = 2053$ in (13). Then

$$\Delta_1(3.1) = \ln(\frac{2}{\sqrt{3.1}}) - \frac{\ln 2053}{2052} > 0.1237\,,$$

and $L_1 = 2125$ can be taken in (17). Checking (10) for $3 \leq l < 2125$, we obtain that $3 \leq l \leq 510$. For all these l such that (10) which is

$$\frac{\varphi(l)}{2} \cdot \left(\ln \frac{2}{\sqrt{3.1}} - \frac{\ln \gamma(l)}{\varphi(l)}\right) < \ln \sqrt{\frac{14^5}{3.1}} - \ln \sin \frac{\pi}{l} \tag{63}$$

satisfies, we obtain

$$[\mathbb{F}_l : \mathbb{Q}] = \frac{\varphi(l)}{2} \leq 99 \tag{64}$$

where 99 is achieved for $l = 199$. Moreover, for all these non-exceptional l we obtain the bound (18) which is

$$N_0 = [\mathbb{K} : \mathbb{F}_l] \leq \left[\frac{\ln \sqrt{\frac{14^5}{3.1}} - \ln \sin \frac{\pi}{l}}{\frac{\varphi(l)}{2} \cdot \left(\ln \frac{2}{\sqrt{3.1}} - \frac{\ln \gamma(l)}{\varphi(l)}\right)} \right], \tag{65}$$

and finally we obtain the bound (19) which is

$$N = [\mathbb{K} : \mathbb{Q}] \le \left[\frac{\ln \sqrt{\frac{14^5}{3.1}} - \ln \sin \frac{\pi}{l}}{\frac{\varphi(l)}{2} \cdot \left(\ln \frac{2}{\sqrt{3.1}} - \frac{\ln \gamma(l)}{\varphi(l)} \right)} \right] \cdot \frac{\varphi(l)}{2}\ . \tag{66}$$

If either l is exceptional, or the right hand side of (66) is more than 99 (this is possible for $3 \le l \le 113$ only), we also apply to l the method A of the Case 1 to improve the poor bound (65) for $N_0 = [\mathbb{K} : \mathbb{F}_l]$ for non-exceptional l. We can apply this method to any $l \ge 4$ since (21) is valid for all $l \ge 4$ if $a_1 + a_2 = 6.2$. For $l \ge 6$, this method gives what we want: $[\mathbb{K} : \mathbb{Q}] \le 99$. For $l = 4$, it only gives $[\mathbb{K} : \mathbb{Q}] \le 120$; for $l = 5$, it only gives $[\mathbb{K} : \mathbb{Q}] \le 172$.

If $l = 3, 4, 5$, equivalently $m = 3, 4, 5$, considering the subgraph of e, δ_4, δ_2, exactly the same consideration as for the graph $\Gamma_1^{(5)}(14)$ above for $m_1 = 3, 4, 5$, give that $[\mathbb{K} : \mathbb{Q}] \le 76$ for $m = 3$, $[\mathbb{K} : \mathbb{Q}] \le 31$ for $m = 4$, and $[\mathbb{K} : \mathbb{Q}] \le 24$ for $m = 5$.

Thus, $[\mathbb{K} : \mathbb{Q}] \le 99$ for all graphs $\Gamma_4^{(6)}(14)$.

This finishes the proof of Theorem 2.7.

4 Appendix: Some results about cyclotomic fields

This is exactly the same as in [20]. We repeat it for readers convenience.

Here we give some results about cyclotomic fields which we used. All of them follow from standard results. For example, see the book [5].

We consider the cyclotomic field $\mathbb{Q}\left(\sqrt[l]{1}\right)$ and its totally real subfield $\mathbb{F}_l = \mathbb{Q}\left(\cos\left(2\pi/l\right)\right)$. We have $[\mathbb{Q}\left(\sqrt[l]{1}\right) : \mathbb{Q}] = \varphi(l)$ where $\varphi(l)$ is the Euler function. We have $\mathbb{F}_l = \mathbb{Q}\left(\sqrt[l]{1}\right) = \mathbb{Q}$ for $l = 1, 2$, and $[\mathbb{F}_l : \mathbb{Q}] = \varphi(l)/2$ for $l \ge 3$. It is known (e.g., see [5]) that the discriminant of the field $\mathbb{Q}\left(\sqrt[l]{1}\right)$ is equal to (where p is prime)

$$|\mathrm{discr}\, \mathbb{Q}(\sqrt[l]{1})| = \frac{l^{\varphi(l)}}{\prod_{p|l} p^{\varphi(l)/(p-1)}}\ . \tag{67}$$

Let $\zeta_l = \exp\left(2\pi i/l\right)$ be a primitive l-th root of 1. The element ζ_l generates the ring of integers of $\mathbb{Q}\left(\sqrt[l]{1}\right)$. Further we assume that $l \ge 3$. The equation of ζ_l over $\mathbb{F}_l$ is $g(x) = (x - \zeta_l)(x - \zeta_l^{-1}) = x^2 - (\zeta_l + \zeta_l^{-1})x + 1 = 0$. We have $g'(\zeta_l) = 2\zeta_l - (\zeta_l + \zeta_l^{-1}) = \zeta_l - \zeta_l^{-1}$. Thus,

$$N_{\mathbb{Q}(\sqrt[l]{1})/\mathbb{F}_l}(g'(\zeta_l)) = (\zeta_l - \zeta_l^{-1})(\zeta_l^{-1} - \zeta_l) = 4\sin^2\left(2\pi/l\right)$$

which gives the discriminant $\delta\left(\mathbb{Q}\left(\sqrt[l]{1}\right)/\mathbb{F}_l\right) = 4\sin^2\left(2\pi/l\right)$. It follows

$$|\delta\left(\mathbb{Q}\left(\sqrt[l]{1}\right)/\mathbb{Q}\right)| = |\delta\left(\mathbb{F}_l/\mathbb{Q}\right)^2 N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2\left(2\pi/l\right))|.$$

We have

$$N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2(\pi/l)) = \gamma(l) = \begin{cases} p & \text{if } l = p^t > 2 \text{ where } p \text{ is prime,} \\ 1 & \text{otherwise.} \end{cases} \tag{68}$$

If l is odd, then $4\sin^2(\pi/l)$ and $4\sin^2(2\pi/l)$ are conjugate, and their norms are equal. Thus,

$$N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2(2\pi/l)) = \gamma(l), \text{ if } l \geq 3 \text{ is odd.}$$

If l is even and $l_1 = l/2$, then $4\sin^2(2\pi/l) = 4\sin^2(\pi/l_1)$. If l_1 is odd, then $\mathbb{F}_{l_1} = \mathbb{F}_l$, and we get

$$N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2(2\pi/l)) = \gamma(l/2) \text{ if } l \geq 6 \text{ is even, but } l/2 \text{ is odd.}$$

If $l_1 \geq 4$ is even, then $[\mathbb{F}_l : \mathbb{F}_{l_1}] = 2$, and we get

$$N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2(2\pi/l)) = \gamma(l/2)^2 \text{ if } l \geq 8 \text{ and } l/2 \text{ is even.}$$

At last,

$$N_{\mathbb{F}_4/\mathbb{Q}}(4\sin^2(2\pi/4)) = 4$$

if $l = 4$.

Thus, finally we get for $l \geq 3$:

$$N_{\mathbb{F}_l/\mathbb{Q}}(4\sin^2(2\pi/l)) = \widetilde{\gamma}(l) = \begin{cases} \gamma(l) & \text{if } l \geq 3 \text{ is odd,} \\ \gamma(l/2) & \text{if } l/2 \geq 3 \text{ is odd,} \\ \gamma(l/2)^2 & \text{if } l/2 \geq 4 \text{ is even,} \\ 4 & \text{if } l = 4. \end{cases} \tag{69}$$

Moreover, we obtain the formula for the discriminant:

$$|\operatorname{discr} \mathbb{F}_l| = \Big(|\operatorname{discr} \mathbb{Q}(\sqrt[l]{1})| \,/\, \widetilde{\gamma}(l) \Big)^{1/2} \quad \text{for } l \geq 3 \tag{70}$$

where $|\operatorname{discr} \mathbb{Q}(\sqrt[l]{1})|$ is given by (67), and $\widetilde{\gamma}(l)$ is given by (69).

We denote $\mathbb{F}_{k,s} = \mathbb{Q}(\cos(2\pi/k), \cos(2\pi/s))$. Further we assume that $k, s \geq 3$. Let $m = [k, s]$ be the least common multiple of k and s. Then $\mathbb{F}_{k,s} \subset \mathbb{F}_m \subset \mathbb{Q}(\sqrt[m]{1})$. We have $\operatorname{Gal}\left(\mathbb{Q}(\sqrt[m]{1})/\mathbb{Q}\right) = (\mathbb{Z}/m\mathbb{Z})^*$ where $\alpha \in (\mathbb{Z}/m\mathbb{Z})^*$ acts on each m-th root ζ of 1 by the formula $\zeta \mapsto \zeta^\alpha$. Obviously, $\mathbb{F}_{k,s}$ is the fixed field of the subgroup G of the Galois group $(\mathbb{Z}/m\mathbb{Z})^*$ which consists of all $\alpha \in (\mathbb{Z}/m\mathbb{Z})$ such that $\alpha \equiv \pm 1 \bmod k$ and $\alpha \equiv \pm 1 \bmod s$. The G includes the subgroup of order two of $\alpha \equiv \pm 1 \bmod m$. If $\alpha \equiv 1 \bmod k$, then $\alpha \equiv 1 + kt \bmod m$ where $t \in \mathbb{Z}$. If $1 + kt \equiv -1 \bmod s$, then the equation $kt + sr = 2$ has an integer solution (t, r) which is equivalent to $(k, s)|2$. Thus, G has the order 4 if and only if $(k, s)|2$. Otherwise, G has the order 2. We set

$$\rho(k, s) = \begin{cases} 2 & \text{if } (k, s)|2, \\ 1 & \text{otherwise,} \end{cases} \tag{71}$$

and we obtain

$$[\mathbb{F}_{k,s} : \mathbb{Q}] = \frac{\varphi(m)}{2\rho(k,s)}. \tag{72}$$

Moreover, we get

$$\mathbb{F}_{k,s} = \mathbb{F}_m \text{ if } (k,s) \not| 2.$$

It follows,

$$|\operatorname{discr} \mathbb{F}_{k,s}| = |\operatorname{discr} \mathbb{F}_m| \text{ if } (k,s) \not| 2, \tag{73}$$

where $m = [k,s]$, and $|\operatorname{discr} \mathbb{F}_m|$ is given by (70).

Assume that $(k,s)|2$. If $(k,s) = 1$, then the fields $\mathbb{Q}(\sqrt[k]{1})$ and $\mathbb{Q}(\sqrt[s]{1})$ are linearly disjoint and their discriminants are coprime. Then their subfields $\mathbb{F}_k$ and $\mathbb{F}_s$ are linearly disjoint, and their discriminants are coprime, and we obtain

$$|\operatorname{discr} \mathbb{F}_{k,s}| = |\operatorname{discr} \mathbb{F}_k|^{(\varphi(s)/2)} |\operatorname{discr} \mathbb{F}_s|^{(\varphi(k)/2)} \text{ if } (k,s) = 1 \text{ and } k,s \geq 3.$$

Assume that $(k,s) = 2$. Then one of $k/2$ or $s/2$ is odd. Assume, $k_1 = k/2$ is odd. Then $\mathbb{F}_k = \mathbb{F}_{k_1}$ and $\mathbb{F}_{k,s} = \mathbb{F}_{k_1,s}$ where $(k_1, s) = 1$. Thus, we obtain the previous case which gives exactly the same formula. We finally obtain

$$|\operatorname{discr} \mathbb{F}_{k,s}| = |\operatorname{discr} \mathbb{F}_k|^{(\varphi(s)/2)} |\operatorname{discr} \mathbb{F}_s|^{(\varphi(k)/2)} \text{ if } (k,s)|2 \text{ and } k,s \geq 3 \tag{74}$$

where the discriminants $|\operatorname{discr} \mathbb{F}_k|$ and $|\operatorname{discr} \mathbb{F}_s|$ are given by (70).

References

[1] I. Agol, *Finiteness of arithmetic Kleinian reflection groups*, Proceedings of the International Congress of Mathematicians, Madrid, 2006, Vol. 2, 951–960 (see also math.GT/0512560).

[2] I. Agol, M. Belolipetsky, P. Storm, K. Whyte, *Finiteness of arithmetic hyperbolic reflection groups*, Preprint 2006, Arxiv:math.GT/0612132, 16 pages.

[3] R.E. Borcherds, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), no. 2, 405–444.

[4] A. Borel, *Commensurability classes and volumes of hyperbolic 3-manifolds*, Ann. Scuola Norm. Sup. Pisa, **8** (1981), 1– 33.

[5] J.W.S. Cassels, A. Fröhlich (eds), *Algebraic number theory*, Proc. of an instructional conference organized by LMS, Academic Press, 1967, xviii+366 pp.

[6] J.H. Conway, J. H., S.P. Norton, *Monstrous moonshine*, Bull. London Math. Soc. **11** (1979), no. 3, 308–339.

[7] H.S.M. Coxeter, *Discrete groups generated by reflections*, Ann. of Math. **35** (1934), no. 2, 588–621.

[8] V.A. Gritsenko, V.V. Nikulin, *Automorphic forms and Lorentzian Kac-Moody algebras. I*, Internat. J. Math. **9** (1998), no. 2, 153–199.

[9] V.A. Gritsenko, V.V. Nikulin, *Automorphic forms and Lorentzian Kac-Moody algebras. II*, Internat. J. Math. **9** (1998), no. 2, 201–275.

[10] V.A. Gritsenko, V.V. Nikulin, *The arithmetic mirror symmetry and Calabi-Yau manifolds*, Comm. Math. Phys. **210** (2000), no. 1, 1–11.

[11] V.A. Gritsenko, V.V. Nikulin, *On the classification of Lorentzian Kac-Moody algebras*, Uspekhi Mat. Nauk **57** (2002) no. 5, 79–138; English transl. in Russian Math. Surveys **57** (2002), no. 5, 921–979.

[12] F. Lannér *On complexes with transitive groups of automorphisms*, Comm. Sém., Math. Univ. Lund **11** (1950), 1–71.

[13] D.D. Long, C. Maclachlan, A.W. Reid, *Arithmetic Fuchsian groups of genus zero*, Pure and Applied Mathematics Quarterly, **2** (2006), no. 2, 1–31.

[14] V.V. Nikulin, *On arithmetic groups generated by reflections in Lobachevsky spaces*, Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), no. 3, 637–669; English transl. in Math. USSR Izv. **16** (1981), no. 3, 573–601.

[15] V.V. Nikulin, *On the classification of arithmetic groups generated by reflections in Lobachevsky spaces*, Izv. Akad. Nauk SSSR Ser. Mat. **45** (1981), no. 1, 113–142; English transl. in Math. USSR Izv. **18** (1982), no. 1, 99–123.

[16] V.V. Nikulin, *Discrete reflection groups in Lobachevsky spaces and algebraic surfaces*, Proceedings of the International Congress of Mathematicians, Berkeley, 1986, Vol. 1, 1987, pp. 654-671.

[17] V.V. Nikulin *The remark on arithmetic groups generated by reflections in Lobachevsky spaces*, Preprint Max-Planck-Institut für Mathematik, Bonn. MPI/91 (1991).

[18] V.V. Nikulin, *Finiteness of the number of arithmetic groups generated by reflections in Lobachevski spaces*, Izvestiya:Mathematics, **71** (2007), no. 1, 53–56 (see also arXiv.org:math.AG/0609256).

[19] V.V. Nikulin, *On ground fields of arithmetic hyperbolic reflection groups*, Preprint arXiv:0708.3991 [math.AG], 33 pages.

[20] V.V. Nikulin, *On ground fields of arithmetic hyperbolic reflection groups.II*, Preprint arXiv:0710.0162 [math.AG], 27 pages.

[21] K. Takeuchi, *A characterization of arithmetic Fuchsian groups,* J. Math. Soc. Japan, **27** (1975), no. 4, 600–612.

[22] K. Takeuchi, *Arithmetic triangle groups,* J. Math. Soc. Japan **29** (1977), no. 1, 91–106.

[23] K. Takeuchi, *Commensurability classes of arithmetic triangle groups,* J. Fac. Sci. Univ. Tokyo Sect. 1A Math. **24** (1977), no. 1, 201-212.

[24] K. Tekeuchi, *Arithmetic Fuchsian groups with signature* $(1, e)$*,* J. Math. Soc. Japan **35** (1983), no. 3, 381–407.

[25] M.-F. Vigneras, *Quelques remarques sur la conjecture* $\lambda_1 \geq \frac{1}{4}$, Seminar on number theory, Paris 1981-82, Progr. Math., **38**, Birkhauser, (1983), 321–343.

[26] É.B. Vinberg, *Discrete groups generated by reflections in Lobačevskiĭ spaces*, Mat. Sb. (N.S.) **72** (1967), 471–488; English transl. in Math. USSR Sb.**1** (1967), 429–444.

[27] É.B. Vinberg, *The nonexistence of crystallographic reflection groups in Lobachevskiĭ spaces of large dimension*, Funkts. Anal. i Prilozhen. **15** (1981), no. 2, 67–68; English transl. in Funct. Anal. Appl. **15** (1981), 216–217.

[28] É.B. Vinberg, *Absence of crystallographic reflection groups in Lobachevskiĭ spaces of large dimension*, Trudy Moskov. Mat. Obshch. **47** (1984), 68–102; English transl. in Trans. Moscow Math. Soc. **47** (1985).

[29] É.B. Vinberg, *Discrete reflection groups in Lobachevsky spaces*, Proceedings of the International Congress of Mathematicians, Warsaw, 1983, Vol. 1, 1984, pp. 593–601.

[30] É.B. Vinberg *Hyperbolic groups of reflections*, Uspekhi Mat. Nauk **40** (1985), no. 1, 29–66; English transl. in Russian. Math. Surv. **40** (1985), no. 1, 31–75.

[31] I.M. Vinogradov, *An introduction to the theory of numbers*, Pergamon, London, 1955.

[32] P. Zograf, *A spectral proof of Rademacher's conjecture for congruence subgroups of the modular group*, J. für die reine agew. Math. **414** (1991), 113–116.

V.V. Nikulin
Deptm. of Pure Mathem. The University of Liverpool, Liverpool
L69 3BX, UK;
Steklov Mathematical Institute,
ul. Gubkina 8, Moscow 117966, GSP-1, Russia
vnikulin@liv.ac.uk   vvnikulin@list.ru